TABLE 4Trends in Prevalence of Use of Smokeless Tobacco for Eighth, Tenth, and Twelfth Graders

10E 104

																							'05–'06
	<u> 1975-85</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>	<u>change</u>
Lifetime																							
8th Grade	_			_	_	—	22.2	20.7	18.7	19.9	20.0	20.4	16.8	15.0	14.4	12.8	11.7	11.2	11.3	11.0	10.1	10.2	+0.1
10th Grade	_	_	_	_	_	_	28.2	26.6	28.1	29.2	27.6	27.4	26.3	22.7	20.4	19.1	19.5	16.9	14.6	13.8	14.5	15.0	+0.6
12th Grade	_	31.4	32.2	30.4	29.2	_	—	32.4	31.0	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	18.3	17.0	16.7	17.5	15.2	-2.3
Thirty-Day																							
8th Grade	_	_		_	_	_	6.9	7.0	6.6	7.7	7.1	7.1	5.5	4.8	4.5	4.2	4.0	3.3	4.1	4.1	3.3	3.7	+0.3
10th Grade	_	_		_	_	_	10.0	9.6	10.4	10.5	9.7	8.6	8.9	7.5	6.5	6.1	6.9	6.1	5.3	4.9	5.6	5.7	+0.1
12th Grade	—	11.5	11.3	10.3	8.4	—	—	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	7.6	6.1	-1.5
Daily																							
8th Grade	_	_	_	_	_	_	1.6	1.8	1.5	1.9	1.2	1.5	1.0	1.0	0.9	0.9	1.2	0.8	0.8	1.0	0.7	0.7	+0.1
10th Grade	_	_		_	_	_	3.3	3.0	3.3	3.0	2.7	2.2	2.2	2.2	1.5	1.9	2.2	1.7	1.8	1.6	1.9	1.7	-0.2
12th Grade	—	4.7	5.1	4.3	3.3	—	—	4.3	3.3	3.9	3.6	3.3	4.4	3.2	2.9	3.2	2.8	2.0	2.2	2.8	2.5	2.2	-0.3
Approx. Ns:																							
8th Grade				_		_	17 500	18 600	18 300	17 300	17,500	17 800	18 600	18 100	16 700	16 700	16 200	15 100	16 500	17 000	16 800	16 500	
10th Grade	_	_				_					17,000												
12th Grade		15 200	14 200	16 200	16 700	15 200	15,000																
								19,800	10,300	19,400	13,400	14,300	19,400	13,200	13,000	12,800	12,800	12,900	14,000	14,000	14,700	14,200	

Source: The Monitoring the Future Study, the University of Michigan.

Notes: Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001.

'--' indicates data not available.

Any apparent inconsistency between the change estimate and the prevalence of use estimates for the two most recent classes is due to rounding error.

For 8th and 10th graders: Data based on one of two forms for 1991–96 and on two of four forms beginning in 1997; N is one-half of N indicated.

For 12th graders: Data based on one form; N is one-fifth of N indicated in 1986–1988 and one-sixth of N indicated beginning in 1989.

The prevalence of use of smokeless tobacco was not asked of 12th graders in 1990 and 1991. Prior to 1990 the prevalence of use question on smokeless tobacco was located near the end of one 12th-grade questionnaire form, whereas after 1991 the question was placed earlier and in a different form. This shift could explain the discontinuities between the corresponding data.